



UNIVERSITI PUTRA MALAYSIA

**APPLICATION OF SEWAGE SLUDGE ON AN ACID TROPICAL SOIL:
CROP RESPONSE AND HEAVY METALS UPTAKE BY MAIZE
AND THEIR ACCUMULATION IN THE SOIL**

ROSAZLIN BINTI ABDULLAH

FP 2002 28

**APPLICATION OF SEWAGE SLUDGE ON AN ACID TROPICAL SOIL:
CROP RESPONSE AND HEAVY METALS UPTAKE BY MAIZE
AND THEIR ACCUMULATION IN THE SOIL**

ROSAZLIN BINTI ABDULLAH

**MASTER OF AGRICULTURAL SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2002



**APPLICATION OF SEWAGE SLUDGE ON AN ACID TROPICAL SOIL:
CROP RESPONSE AND HEAVY METALS UPTAKE BY MAIZE
AND THEIR ACCUMULATION IN THE SOIL**

By

ROSAZLIN BINTI ABDULLAH

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfillment of the Requirements for the
Degree of Master of Agricultural Science**

July 2002



DEDICATION

This thesis is dedicated to:

My beloved parents

Abdullah Md Noh

and

Junidah Hamid

Brother and sisters,

Rozaidey and Norhayati

Rosazura

Rosazlinda

Rosazleza

Rosazliyana

Who always supported and encouraged me to do the best

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Agricultural Science

**APPLICATION OF SEWAGE SLUDGE ON AN ACID TROPICAL SOIL:
CROP RESPONSE AND HEAVY METALS UPTAKE BY MAIZE
AND THEIR ACCUMULATION IN THE SOIL**

By

ROSAZLIN BINTI ABDULLAH

July 2002

Chairman : Dr. Che Fauziah Binti Ishak

Faculty : Agriculture

At present, Malaysia produces about 5 million m³ sludge per year (wet weight basis). Land application of sewage sludge is considered the most economical and also provides an opportunity to recycle beneficial plant nutrients and organic matter to soil for crop production. However, sewage sludge also contain varying amounts of heavy metals which may pose a hazard of metal toxicity to crops and the consumer of these crops. Thus the uptake of heavy metals by crops and the fate of these heavy metals in soils need to be monitored.

The first study, which involved ten sewage sludges, were collected from different wastewater treatment plants and classified into domestic sludge (type A) and light-industry sludge mixed with domestic sludge (type B). The chemical properties of sewage sludge with emphasis on heavy metal content were determined. In general, type

B sludge has higher heavy metal content than type A sludge. The concentration of heavy metals in these sludges did not exceed the maximum permitted concentrations (MPC) of the European Community Standard (ECS), for land application, except for Zn (7110.10 mg kg⁻¹). Also, the Cu concentration is rather high in type B sludge. Therefore, type B sludge can be characterized as quite hazardous and unsuitable for agricultural use.

The second study was a field experiment conducted at the share farm Universiti Putra Malaysia from 1999 till 2001. The study was to investigate the crop response and uptake of heavy metals from sewage sludge by maize, and, to correlate heavy metals in the soil with content in the maize. The treatments carried out were inorganic N (ammonium sulfate) at 140 kg N ha⁻¹, control (no N application) and sludge application rates at 140, 280, 420 and 560 kg N ha⁻¹ equivalent with five replications. Application of sewage sludge and inorganic fertilizers produced significantly higher yield than the control. The rate of 420 kg ha⁻¹ sewage sludge gave the highest dry matter yield for the 1st cycle and 720 kg N ha⁻¹ sewage sludge for the 3rd cycle. It can be concluded that sewage sludge was able to perform just as good as inorganic fertilizer. Sewage sludge application seemed to have little effect in increasing heavy metals concentrations in the soil and grain after the third maize cycle. The concentration of heavy metals in the soil were below the MPC of the ECS, which ranged: Zn (12.44-35.44 mg kg⁻¹), Cu (5.00-9.80 mg kg⁻¹), Cd (0.60-2.44 mg kg⁻¹), Pb (7.16-24.04 mg kg⁻¹), Ni (7.44-11.36 mg kg⁻¹), Mn (29.76-41.32 mg kg⁻¹), Cr (19.72-44.00 mg kg⁻¹) and Fe (1.47-2.23 %), respectively. The concentrations of heavy metals in maize grain were below the MPC values of Malaysian Food Act 1983 and Food Regulation 1985 (fresh weight basis) which ranged : Zn (4.95-19.18 mg

kg⁻¹), Cu (0.56-2.60 mg kg⁻¹), Cd (0.037-0.052 mg kg⁻¹), Pb (0.034-0.052 mg kg⁻¹), Ni (0.66-1.22 mg kg⁻¹), Mn (1.56-8.53 mg kg⁻¹), Cr (0.12-0.44 mg kg⁻¹) and Fe (8.16-24.93 mg kg⁻¹), respectively. There were significant correlations between total heavy metals in soil and content of heavy metals in leaves and stems.

The third study is to determine the forms of Cd, Cu, Ni, Pb and Zn in the sewage sludge, and in the soils applied with sewage sludge, after the 1st and 3rd maize cycle. The correlations between different forms of heavy metals in the soil and content in maize grain were also investigated. Fractionation of heavy metals in sewage sludge showed that the dominant form of all heavy metals were residual form except Cu. Leaving the residual fraction out, Cd and Pb were dominant in exchangeable form, Ni in carbonate form and Zn in Fe-Mn oxide form. The organic form is dominant for Cu in sludge treated soil. Cd and Pb were predominant in the exchangeable form. Ni was predominant in the carbonate form and Zn was predominant in the Fe-Mn oxide form. In general, the percentage of water soluble content was less than 5%. Significant correlations were only obtained between Cd content in maize grain and the organic forms in soil, Ni content in grain with total metal in the soil and between Cu content in maize grain and the carbonate, Fe-Mn oxide and organic forms in soils.

For conclusion, domestic sewage sludge produced in Malaysia can be safely as fertilizer applied on agricultural land, in three continuous applications, for crop production. However, the Cu and Zn concentrations in grain and soil need to be monitored.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia bagi memenuhi keperluan penganugerahan ijazah Master Sains Pertanian

**PENGUNAAN ENAP CEMAR KUMBAHAN PADA TANAH ASID
TROIKA: HASIL PERTUMBUHAN DAN PENGAMBILAN LOGAM
BERAT OLEH JAGUNG DAN PENGUMPULANNYA DI DALAM TANAH**

Oleh

ROSAZLIN BINTI ABDULLAH

Julai 2002

Pengerusi : Dr. Che Fauziah Binti Ishak

Fakulti : Pertanian

Setiap tahun Malaysia menghasilkan lebih daripada 5 juta tan enap cemar kumbahan (berdasarkan berat basah). Penggunaan enap cemar kumbahan tersebut pada tanah dapat mengurangkan kos dan ia juga mengandungi nutrien dan bahan organik untuk tanaman. Walaubagaimanapun, enap cemar kumbahan juga mengandungi logam berat yang menyebabkan ketoksikan kepada tanaman dan kepada pengguna tanaman. Jadi, pemantauan pengambilan logam berat oleh tanaman dan tindakbalas logam berat dalam tanah perlu dilakukan.

Kajian pertama melibatkan sepuluh jenis enap cemar kumbahan yang diambil daripada pelbagai loji rawatan dan dibahagikan kepada enap cemar domestik (jenis A) dan enap cemar industri ringan bercampur dengan enap cemar domestik (jenis B). Komposisi kimia enap cemar kumbahan dengan penekanan kepada kandungan logam

berat dikenalpasti. Pada umumnya, enap cemar kumbahan jenis B adalah lebih tinggi berbanding kandungan logam berat enap cemar jenis A. Kepekatan logam berat di dalam enap cemar kumbahan tidak melebihi tahap yang dibenarkan (MPC) oleh garis panduan Kesatuan Eropah (ECS) untuk penggunaannya ke atas tanah kecuali Zn (7110 mg kg^{-1}). Juga, kepekatan Cu yang agak tinggi di dalam enap cemar kumbahan jenis B. Oleh itu, enap cemar jenis boleh diklasifikasi agak merbahaya dan tidak sesuai untuk digunakan pada tanah pertanian.

Kajian kedua ialah kajian di ladang yang telah dijalankan di Ladang Kongsi Universiti Putra Malaysia dari tahun 1999 hingga tahun 2001. Kajian ini adalah untuk mengenalpasti kesan pertumbuhan oleh tanaman dan pengambilan logam berat daripada enap cemar kumbahan oleh jagung, dan, korelasi antara logam berat di dalam tanah dengan kandungannya di dalam jagung. Rawatan yang digunakan ialah baja tak organik N (ammonium sulfat) pada kadar 140 kg N ha^{-1} , kawalan (tiada penggunaan N), dan penggunaan enap cemar kumbahan pada kadar setara 140, 280, 420 dan 560 kg N ha^{-1} dengan 5 replikasi. Penggunaan enap cemar kumbahan dan baja tak organik menunjukkan hasil yang tinggi berbanding kawalan. Kadar enap cemar 420 kg N ha^{-1} memberikan hasil yang tertinggi pada tanaman pertama dan 720 kg N ha^{-1} enap cemar pada tanaman ketiga. Ini boleh disimpulkan bahawa enap cemar kumbahan mampu membekal N seperti baja tak organik. Penggunaan enap cemar kumbahan ke atas tanaman memberi kesan ke atas tanah dan bijirin selepas tiga kali penanaman. Kepekatan logam berat dalam tanah masih di bawah tahap yang dibenarkan (MPC) oleh ECS iaitu : Zn ($12.44\text{--}35.44 \text{ mg kg}^{-1}$), Cu ($5.00\text{--}9.80 \text{ mg kg}^{-1}$), Cd ($0.60\text{--}2.44 \text{ mg kg}^{-1}$), Pb ($7.16\text{--}24.04 \text{ mg kg}^{-1}$), Ni ($7.44\text{--}11.36 \text{ mg kg}^{-1}$), Mn

(29.76-41.32 mg kg⁻¹), Cr (19.72-44.00 mg kg⁻¹) dan Fe (1.47-2.23 %). Kepekatan logam berat di dalam biji jagung masih di bawah tahap yang dibenarkan (MPC) menurut Akta Makanan Malaysia 1983 dan Peraturan Makanan 1985 (berdasarkan berat basah) iaitu :Zn (4.95-19.18 mg kg⁻¹), Cu (0.56-2.60 mg kg⁻¹), Cd (0.037-0.052 mg kg⁻¹), Pb (0.034-0.052 mg kg⁻¹), Ni (0.66-1.22 mg kg⁻¹), Mn (1.56-8.53 mg kg⁻¹), Cr (0.12-0.44 mg kg⁻¹) dan Fe (8.16-24.93 mg kg⁻¹). Terdapat perkaitan yang signifikan di antara logam berat di dalam tanah dan kandungan logam berat di dalam daun dan batang.

Kajian ketiga ialah untuk mengenalpasti bentuk Cd, Cu, Ni, Pb dan Zn di dalam enap cemar kumbahan dan juga di dalam tanah yang dirawat oleh enap cemar kumbahan selepas tanaman pertama dan ketiga. Korelasi di antara bentuk logam berat yang berbeza dengan kandungannya di dalam biji jagung juga dikaji. Kajian pemerinkatan di dalam enap cemar kumbahan menunjukkan semua logam berat yang dikaji adalah dominan di dalam bentuk sisa baki kecuali Cu. Jika bentuk sisa baki tidak di ambil kira, Cd dan Pb adalah dominan dalam bentuk tukarganti, Ni dominan dalam bentuk karbonat dan Zn dalam bentuk Fe-Mn oksida. Cu adalah dominan dalam bentuk organik bagi tanah yang di rawat dengan enap cemar kumbahan, Cd dan Pb adalah dominan dalam bentuk tukarganti, Ni dominan dalam bentuk karbonat, dan Zn dalam bentuk Fe-Mn oksida. Pada umumnya, didapati peratus bentuk larut air yang diekstrak dengan air suling adalah kurang daripada 5%. Terdapat hanya perkaitan positif di antara kandungan Cd di dalam biji jagung dengan bentuk organik di dalam tanah, kandungan Ni di dalam biji jagung dengan bentuk

jumlah di dalam tanah dan di antara kandungan Cu di dalam biji jagung dengan bentuk karbonat, bentuk Fe-Mn oksida dan bentuk organik di dalam tanah.

Kesimpulannya, enap cemar kumbahan domestik yang dihasilkan di Malaysia adalah selamat apabila digunakan sebagai baja untuk tanah pertanian setelah tiga penanaman yang berterusan. Walaubagaimanapun, kepekatan Zn dan Cu dalam bijirin dan tanah hendaklah sentiasa dipantau.

ACKNOWLEDGEMENTS

Alhamdulillah, first of all I would like to express my utmost thanks and gratitude to Almighty Allah SWT Who has given me the will and strength to complete this project and *salawat* and *salam* to His messenger, Prophet Muhammad SAW.

I would also like to express my most sincere appreciation and deepest gratitude to Dr. Che Fauziah Ishak, the chairman of my Supervisory Committee, for her kind assistance, advice and guidance during the course of this study and in the preparation of this thesis.

I am also grateful to other members of the Supervisory Committee, Associate Professor Dr. Rosenani Abu Bakar and Associate Professor Dr. Siti Zauyah Darus, for their supervision, comments and direct contribution to this study.

I am also grateful to the laboratory staffs of the Department of Land Management, Faculty of Agriculture, UPM, especially Puan Norhashimah, Puan Rozita, Cik Norazlina Shaari, En. Rahim, En. Jamil, En. Alias, Mr. Mutuviren, Puan Faridah, Hajah Faridah, Hajah Umi Kalthum, Puan Sarimah, Puan Rusnah, En. Ariffin, Puan Fauziah, and En. Ramli. To my friends, Cik Sharol Nasidah, En. Tai, En. Sabri, Cik Susilawati and Cik Wan Asrina, my sincere thanks for their help and co-operation during the course of my study.

I wish to express my heartfelt thanks to my friends, Cik Aniza Meor Sleman for her patience, support and understanding. As for Cik Noor Hazlyna Din, heartiest thanks for her encouragements and support.

Finally, I would also like to acknowledge the financial support provided by Indah Water Konsortium, which made my study possible.

I certify that an Examination Committee met on 22nd July 2002 to conduct the final examination of Rosazlin Binti Abdullah on her Master of Agricultural Science thesis entitled “Application of Sewage Sludge on an Acid Tropical Soil: Crop Response and Heavy Metals Uptake by Maize and Their Accumulation in the Soil” in accordance with Universiti Putra Malaysia (Higher Degree) Act 1980 and Universiti Putra Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

HALIMI BIN MOHD SAUD, Ph.D.

Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

CHE FAUZIAH BINTI ISHAK, Ph.D.

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROSENANI BINTI ABU BAKAR, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

SITI ZAUYAH BINTI DARUS, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

SHAMSHER MOHAMAD RAMADILI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Master of Agricultural Science. The members of the Supervisory Committee are as follow:

CHE FAUZIAH BINTI ISHAK, Ph.D.

Faculty of Agriculture
Universiti Putra Malaysia
(Chairperson)

ROSENANI BINTI ABU BAKAR, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

SITI ZAUYAH BINTI DARUS, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

AINI IDERIS, Ph.D.

Professor/Dean,
School of Graduate Studies
Universiti Putra Malaysia

Date:

I hereby declare the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



ROSAZLIN BINTI ABDULLAH

Date: 19/09/02

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	vi
APPROVAL SHEETS	x
DECLARATION FORM	xiv
LIST OF TABLES	xix
LIST OF FIGURES	xx
LIST OF PLATES	xxii
LIST OF ABBREVIATIONS	xxiii
 CHAPTER	
 I INTRODUCTION	 1
 II LITERATURE REVIEW	 6
2.1 Heavy Metals	6
2.1.1 Factors Affecting the Bioavailability of Heavy Metal for Plant Uptake	6
2.1.1.1 Soil pH	7
2.1.1.2 Organic Matter	7
2.1.1.3 Cation Exchange Capacity (CEC)	8
2.1.1.4 Plant Factors	9
2.1.1.5 Fertilizer Application	10
2.1.1.6 Environmental Factors	11
2.1.2 Heavy Metals in Sludge	11
2.1.2.1 Loading Limits of Heavy Metals	12
2.1.2.2 Potentially Toxic Elements	16
2.2 Sewage Sludge	19
2.2.1 Sewage Sludge Properties	22
2.2.1.1 Physical Properties	22
2.2.1.2 Chemical Properties	22
2.2.1.3 Biological Properties	23
2.2.2 Land Application of Sewage Sludge	24
2.2.3 Benefits of Sewage Sludge Application on Land	25
2.2.3.1 Sewage Sludge as N Fertilizers	28
2.2.3.2 Nitrogen Mineralization of Sewage Sludge	29
2.2.3.3 Factors Affecting N Mineralization of Sludge	30
2.2.4 Crop Response to Sewage Sludge Application	33
2.2.4.1 Effect of Sludge Properties on Accumulation of Heavy Metals by Crops	35

2.2.4.2	Soil Properties Which Influence the Accumulation of Heavy Metals by Crops	35
2.3	Heavy Metals Fractionation	37
2.3.1	Analytical Considerations	42
III	GENERAL MATERIALS AND METHODS	45
3.1	Preparations and Analyses of Sewage Sludge	45
3.1.1	Determination of Moisture content (based on wet weight) and Total Dry Solid	46
3.1.2	Determination pH of Sewage Sludge	46
3.1.3	Determination of Volatile Solid and Organic Carbon of Sewage Sludge (Loss on Ignition Method)	47
3.1.4	Determination of Total Nitrogen (Kjedahl Method)	48
3.1.5	Determination of Total Heavy Metals in Sewage Sludge	48
3.2	Preparations and Analyses of Soil Samples	49
3.2.1	Determination of Soil pH	49
3.2.2	Determination of Electrical Conductivity (EC)	49
3.2.3	Determination of Organic Carbon (Walkley and Black)	50
3.2.4	Determination of Available Heavy Metals	50
3.2.5	Determination of Available Phosphorous	51
3.2.6	Determination of Cation Exchange Capacity (CEC)	51
3.2.7	Determination of Total Heavy Metals in Soil	52
3.2.8	Determination of Particle Size Distribution (texture)	52
3.3	Preparations and Analyses of Plant Samples	52
3.3.1	Determination of Total Heavy Metals (Dry Ashing Method)	53
3.3.2	Determination Total Nitrogen (Kjedahl Method)	54
IV	STUDY I: CHEMICAL CHARACTERIZATION OF SEWAGE SLUDGE FROM SEVERAL WASTEWATER TREATMENT PLANTS	55
4.1	Introduction	55
4.2	Materials and Methods	56
4.2.1	Sludge Sampling	58
4.2.2	Physical and Chemical Analyses of Sewage Sludge	58
4.2.3	Statistical Analyses	58
4.3	Results and Discussion	59
4.3.1	Physical Properties of Sewage Sludge	63
4.3.2	Chemical Properties Sewage Sludge	64
4.3.2.1	Major/Macro Elements	64
4.3.2.2	Heavy Metals	67
4.4	Conclusion	69

V	STUDY 2: EVALUATION OF SEWAGE SLUDGE AS A SOURCE OF N FERTILIZERS IN THE MAIZE PRODUCTION	71
5.1	Introduction	71
5.2	Materials and Methods	73
5.2.1	Site, Soil and Climate	73
5.2.2	Experimental Treatments and Design	74
5.2.3	Application of Sewage Sludge	78
5.2.4	Sowing	78
5.2.5	Crop Maintenance	80
5.2.6	Harvesting	81
5.2.7	Plant sampling	82
5.2.8	Soil Sampling, Preparation and Analyses	82
5.2.9	Quality Assurance of Cd and Pb Analyses in Grains	83
5.2.10	Statistical Analyses	83
5.3	Results and Discussion	84
5.3.1	Dry Matter Yield	84
5.3.1.1	Field Spatial Variability	87
5.3.1.2	Dry Matter Yield Calibration Curve	90
5.3.2	Economic Yield	92
5.3.3	Effect of Soil Chemical and Physical Properties Applied with Sewage Sludge	93
5.3.3.1	pH	93
5.3.3.2	Electrical Conductivity (EC)	94
5.3.3.3	Total N Content	95
5.3.3.4	Available P	97
5.3.3.5	Organic Carbon Content	98
5.3.3.6	Cation Exchange Capacity (CEC)	99
5.3.4	Heavy Metals in the Soil	100
5.3.5	Heavy Metals in the Maize Grain	108
5.3.6	Heavy Metals in Other Parts of the Maize Plants	116
5.3.7	Nitrogen in Plant Tissue	117
5.3.8	Correlation Study	120
5.3.9	Quality Assurance	122
5.4	Conclusion	123
5.5	Recommendation for Future Studies	125
VI	STUDY 3: FRACTIONATION STUDY OF HEAVY METALS IN SOILS APPLIED WITH SEWAGE SLUDGE	126
6.1	Introduction	126
6.2	Materials and Methods	127
6.2.1	Sewage Sludge and Soil Samples	127
6.2.2	Preparation of Soil Samples	128
6.2.3	Chemical Analysis	128
6.2.3.1	Water Soluble Fraction	129
6.2.3.2	Exchangeable Fraction	129
6.2.3.3	Carbonate Fraction	131
6.2.3.4	Fe-Mn Oxides Fraction	131

6.2.3.5	Organic Fraction	131
6.2.3.6	Residual Fraction	132
6.2.4	Washing Procedure	132
6.2.5	Total Metal Extraction	133
6.2.6	Heavy Metals content in Maize Grain	133
6.2.7	Statistical Analysis	133
6.3	Results and Discussion	134
6.3.1	Forms of Heavy Metals in Sewage Sludge	134
6.3.2	Forms of Heavy Metals in Soil Applied with Sewage Sludge	137
6.3.2.1	Cadmium	137
6.3.2.2	Copper	141
6.3.2.3	Nickel	145
6.3.2.4	Lead	148
6.3.2.5	Zinc	151
6.3.3	Correlation Study	155
6.4	Conclusions	158
VII	GENERAL CONCLUSION	161
	REFERENCES	165
	APPENDICES	181
A)	Soil Profile	181
B)	Raw Data : Study 2	184
C)	Statistics : Study 2	194
D)	Plates : Study 2	211
VITA		225

LIST OF TABLES

Table		Page
2.1	Limitations imposed on heavy metals in sludge application to land	14
2.2	Maximum values for metal concentration in sewage sludge in agriculture, their rate of application and concentrations in sludge treated soil in the Commission of the European Communities directive (Commissions of the European Communities, 1986)....	15
2.3	Effects of heavy metals on human health.....	17
2.4	Sources of heavy metals in domestic wastewater by product type	18
2.5	Yields response to sludge and commercial fertilizer	34
4.1	Location of sewage sludge used in this study	57
4.2	Chemical characterization of domestic sludge (Type A).....	60
4.3	Chemical characterization of light industries sludge mixed with and domestic mixed sludge (Type B).....	61
4.4	T-test comparison with domestic sludge (Type A) and light industries mixed domestic sludge (Type B).	62
5.1	Physico-chemical properties of the soil at the experimental site before planting and before liming.....	76
5.2	Chemical composition of sewage sludge.....	79
5.3	Schedule of planting of maize for three cycles.....	81
5.4	Mean dry matter yield of the 3 crop cycles (kg ha^{-1}).....	86
5.5	Economic yields (kg ha^{-1}) of the 3 crop cycles	93
5.6	Correlation between total and available heavy metal with content in plant parts (n=60).....	121
5.7	Cross-checking data of maize grain analyses obtained from UPM and Rothamsted Laboratories using graphite method.....	122
6.1	Effect of 1 st and 3 rd cycle on the correlation (r) between extractable metals fraction and content in the maize grain	157

LIST OF FIGURES

Figure		Page
5.1	Rainfall distribution during the 3 crop cycles a) 1 st maize cycle in 1999, b) 2 nd maize cycle in 2000 and c) 3 rd cycle in 2000.....	75
5.2	Layout of experimental plots (microplots were established in all replicates of the treatment).....	77
5.3	Effect of sludge rate on the total dry matter yield a) 1 st maize cycle and b) 3 rd maize cycle	91
5.4	Effect of pH in soil applied with sewage sludge.....	94
5.5	Effect of electrical conductivity (EC) in soil applied with sewage sludge.....	95
5.6	Effect of total N in soil applied with sewage sludge.....	96
5.7	Effect of Available P in soil applied with sewage sludge.....	97
5.8	Effect of organic carbon in soil applied with sewage sludge.....	98
5.9	Effect of cation exchangeable capacity (CEC) pH in soil applied with sewage sludge.....	100
5.10	Mean concentration of heavy metals in soil a) Zn b) Cu.....	107
5.10	(cont.) Mean concentration of heavy metals in soil c) Cd d) Pb.....	105
5.10	(cont.) Mean concentration of heavy metals in soil e) Ni f) Mn.....	106
5.10	(cont.) Mean concentration of heavy metals in soil g) Fe and h) Cr ...	107
5.11	Mean concentration of heavy metals in maize grain a) Zn b) Cu.....	112
5.11	(cont.) Mean concentration of heavy metals in maize grain c) Cd d) Pb.....	113
5.11	(cont.) Mean concentration of heavy metals in maize grain e) Ni f) Mn.....	114
5.11	(cont.) Mean concentration of heavy metals in maize grain g) Fe and h) Cr	115
5.12	N yield in different parts of plant a) 1 st maize cycle.....	118

5.12	(cont.) N yield in different parts of plant b) 2 nd maize cycle and c) 3 rd maize cycle	119
6.1	Schematic diagram for the sequential extraction of the different forms of Cd, Cu, Ni, Pb and Zn in soil applied with sewage sludge.....	130
6.2	Forms of Ni, Cd, Zn, Pb and Cu in sewage sludge.....	135
6.3	Forms of cadmium in soil after 1 st (C1) and 3 rd (C3) maize cycle.....	138
6.4	Mean concentration of cadmium in soil applied with sewage sludge after a) 1 st maize cycle and b) 3 rd maize cycle using different chemical extractants	140
6.5	Forms of copper in soil after 1 st and 3 rd maize cycle.....	142
6.6	Mean concentration of copper in soil applied with sewage sludge after a) 1 st maize cycle and b) 3 rd maize cycle using different chemical extractants	144
6.7	Forms of nickel in soil after 1 st and 3 rd maize cycle.....	146
6.8	Mean concentration of nickel in soil applied with sewage sludge after a) 1 st maize cycle and b) 3 rd maize cycle using different chemical extractants	147
6.9	Forms of lead in soil after 1 st and 3 rd maize cycle.....	149
6.10	Mean concentration of lead in soil applied with sewage sludge after a) 1 st maize cycle and b) 3 rd maize cycle using different chemical extractants	150
6.11	Forms of zinc in soil after 1 st and 3 rd maize cycle.....	152
6.12	Mean concentration of zinc in soil applied with sewage sludge after a) 1 st maize cycle and b) 3 rd maize cycle using different chemical extractants	154

LIST OF PLATES

Plate		Page
2.1	The scheduled desludging of individual septic tank by Indah Water's service (Indah Water, 1997).....	21
5.1	Sowing of maize seeds	80
5.2	Soil profile at the right-end of the field	88
5.3	Soil profile at the left-end of the field.....	89
5.4	Comparison of roots of maize from right-end (R1 T5 and R4T5) and left-end (R3 T5 and R5T5) of the field	90

LIST OF ABBREVIATIONS

C	=	Carbon
CEC	=	Cation Exchange Capacity
EC	=	Electrical Conductivity
ECS	=	European Communities Standard
IWK	=	Indah Water Konsortium
MPC	=	Maximum Permitted Concentration
N	=	Nitrogen
P	=	Phosphorous
UK	=	United Kingdom
UPM	=	Universiti Putra Malaysia
USDA	=	United State Department of Agriculture
USEPA	=	United States Environmental Protection Agency